Help Control Mosquitoes that Spread Dengue, Chikungunya, and Zika Viruses





Aside from being itchy and annoying, the bite of an infected female mosquito (*Aedes aegypti or Aedes albopictus*) can spread dengue, chikungunya, or Zika viruses. People become infected with dengue, chikungunya, or Zika after being bitten by an infected mosquito.

- Female mosquitoes lay several hundred eggs on the walls of waterfilled containers. Eggs stick to containers like glue and remain attached until they are scrubbed off. When water covers the eggs, they hatch and become adults in about a week.
- Adult mosquitoes live inside and outside.
- They prefer to bite during the day.
- A few infected mosquitoes can produce large outbreaks in a community and put your family at risk of becoming sick.

Protect Yourself, Your Family, and Community from Mosquitoes



Eliminate standing water in and around your home:

- Once a week, empty and scrub, turn over, cover, or throw out items that hold water, such as tires, buckets, planters, toys, pools, birdbaths, flowerpots, or trash containers. Check inside and outside your home.
- Tightly cover water storage containers (buckets, cisterns, rain barrels) so that mosquitoes cannot get inside to lay eggs.
- For containers without lids, use wire mesh with holes smaller than an adult mosquito.



If you have a septic tank, follow these steps:

- Repair cracks or gaps.
- Cover open vent or plumbing pipes. Use wire mesh with holes smaller than an adult mosquito.



Keep mosquitoes out of your home:

- Use screens on windows and doors.
- Repair holes in screens.
- Use air conditioning when available.



Put plants in soil, not in water.



Drain water from pools when not in use.



Recycle used tires or keep them protected from rain.



Drain & dump any standing water



Weekly, scrub vases & containers to remove mosquito eggs.





Prevent mosquito bites:

 Use an Environmental Protection Agency (EPA)-registered insect repellent with one of the following active ingredients. All EPA-registered insect repellents are evaluated to make sure they are safe and effective.



Keep rain barrels covered tightly.



Active ingredient

Higher percentages of active ingredient provide longer protection

DEET

Picaridin (known as KBR 3023 and icaridin outside the US)

IR3535

Oil of lemon eucalyptus (OLE) or para-menthane-diol (PMD)

2-undecanone



Find the insect repellent that's right for you by using **EPA's search tool***.



Weekly, empty standing water from fountains and bird baths.

* The EPA's search tool is available at: www.epa.gov/insect-repellents/find-insect-repellent-right-you

- Always follow the product label instructions.
- Reapply insect repellent every few hours, depending on which product and strength you choose.
- Do not spray repellent on the skin under clothing.
- If you are also using sunscreen, apply sunscreen first and insect repellent second.
- Treat clothing and gear (such as boots, pants, socks, and tents) with permethrin or purchase permethrin-treated clothing and gear.
 - Treated clothing remains protective after multiple washings. See product information to find out how long the protection will last.
 - If treating items yourself, follow the product instructions carefully.
 - Do **not** use permethrin products, intended to treat clothing, directly on skin.
- Wear long-sleeved shirts and long pants.



Keep septic tanks sealed.



Install or repair window & door screens

For more information, visit:

www.cdc.gov/dengue, www.cdc.gov/chikungunya, www.cdc.gov/zika